



FAMACHA[®]

Information Guide

Originally compiled by the Faculty of Veterinary Science, University of Pretoria, the Onderstepoort Veterinary Institute, the Worm Workshop of the South African Veterinary Association, and the Intervet South Africa, with the support of the Food and Agriculture Organization of the United Nations, the National Wool Growers' Association and the National and Provincial Departments of Agriculture in South Africa.

Modified by Dr. Ray M. Kaplan and Dr. James E. Miller within the framework of USDA SARE grant # LS02-143 to address the use of FAMACHA[®] in the United States.

IMPORTANT NOTE FOR ALL USERS OF FAMACHA[®]:

To properly implement FAMACHA[®], IT IS ESSENTIAL FOR ALL USERS TO:

- Obtain practical hands-on instruction in the use of FAMACHA[®]
- To be sure to understand the information supplied
- Read and carefully follow all the instructions in this guide

WARNING!

As this Information Guide is used in circumstances outside the compilers' and distributors' control, users must undertake to use it at their own risk. The compilers and distributors, and/or any of their employees do not accept liability for any damage or loss suffered by any person because of or arising from the use of this guide.

WHY THE FAMACHA[®] SYSTEM WAS DEVELOPED:

- *Haemonchus contortus* (barber pole worm) is usually the biggest disease problem of sheep and goats throughout the warm regions of the world, particularly in the subtropical and tropical regions. Major production losses and deaths arise where the parasite (worm) is not adequately controlled.
- Due to the overuse of dewormers over many years, resistance to the dewormers is an ever-increasing problem. On many farms in many countries there is a resistance to all the groups of

deworming drugs, and the viability of sheep and goat farming is threatened. No one can rely on the excessive use of drugs alone to control this parasite in the future.

- While most sheep and goats (especially adults) can withstand the unfavorable effects of *Haemonchus*, a small minority cannot.
- Selectively deworming only those animals that require treatment greatly decreases the development of resistance because the eggs produced by the few resistant worms that survive treatment will be greatly diluted by all the eggs produced by the animals that did not receive treatment. In contrast, when all animals are treated and moved to a parasite- “safe,” or “clean” pasture, only resistant worms that survive treatment will produce all the eggs that form the next generation of worms.
- Both resistance (the ability to prevent or suppress infection) and resilience (the ability to withstand the effect of parasites) have been shown to be moderately heritable. This means that sheep and goats can be either culled or selected for these traits.
- Once sheep and goats that are unable to cope with existing worm challenge infections are identified, they can be targeted for special attention without the whole herd or flock having to be treated. In the long term, by culling animals that are repeatedly identified as unable to cope with moderate worm burdens, a more resistant and resilient flock genetically suited to the environment can be bred.

CLINICAL DIAGNOSIS OF ANEMIA: PRINCIPLE ON WHICH FAMACHA® IS BASED.

Blood consists of a clear, fluid part (plasma) and a cellular part (mainly red blood cells). The proportion of red cells to plasma determines whether the animal is healthy or unhealthy. This proportion can be estimated accurately by assessing the color of the mucous membranes of (especially) the eyes. As *Haemonchus* are blood feeders, the effects of a heavy parasite burden in non-resilient animals will therefore be evident as a low ratio of red cells to plasma. This is seen in the mucous membranes of the eyes as a visible paleness generally known as anemia. By monitoring anemia, resilient and susceptible animals can be identified.

USES AND ADVANTAGES:

- A significant drop in the amount and frequency of deworming can be expected for most of the herd or flock, which will reduce the amount of money spent on drugs.
- Because fewer animals are treated, the development of resistance in worm populations will be slowed down.
- In the long term, elimination of non-resilient animals will allow for the breeding of better adapted animals.
- There will probably only be a small to moderate number of sheep or goats that need to be treated at each examination.
- These animals can be treated before the symptoms and effects of anemia become too severe if the flock is examined regularly.
- Individual animals that repeatedly fail to cope with *Haemonchus* in spite of an effectively designed control program can be identified and eliminated from the herd or flock.
- Animals that escaped treatments or were underdosed or improperly drenched (e.g., owing to faulty drenching syringe), can be identified before severe problems occur.

- If an ineffective dewormer for *Haemonchus* is used, this will become apparent because pale mucous membranes should become noticeably red in color within a week or so. Provided protein intake is sufficient and body condition is adequate.
- If there is a severe build-up of infective larvae in the pasture, an early warning of the impending danger can be a sudden increase in the number of anemic animals.
- Paddocks, pens, and pasture that repeatedly present problems can be identified and appropriate action taken.
- The process of inspecting the eyes is quick and can readily be integrated with other activities like vaccination, weighing, condition scoring or counting. In South Africa it is reported that up to 500 sheep can be inspected per hour with good facilities and practice.
- Because animals are examined frequently, other unrelated problems are quickly discovered.
- The technique is very easy and sufficiently reliable once learned under the guidance of a competent instructor.
- Animals become tamer and easier to handle.

PRECAUTIONS AND POTENTIAL PROBLEMS:

- The FAMACHA[®] system should be used only after it had been fully explained and demonstrated by properly trained instructors.
- Only *Haemonchus* infection can be monitored by this technique.
- FAMACHA[®] is only a component of a good management program for *Haemonchus* and cannot be used on its own.
- Other worms (nematodes) are also important. *Trichostrongylus* (Bankrupt worm) is found in sheep and goats throughout the US. *Teladorsagia* (Brown stomach worm) is common in the northern parts of the US. A program for controlling these and other worms may be needed as well. If either of these worms are the primary problem and *Haemonchus* is only present in low to moderate numbers, then FAMACHA[®] may fail to provide a sound basis for treatment decisions.
- Herd or flock fecal egg counts should be monitored on a periodic basis.
- Animals should be monitored regularly (at least every 2-3 weeks during the *Haemonchus* transmission season, and possibly as often as weekly at the peak of the worm season).
- Animals should always be scored with the help of the chart, not from memory.
- Kids/Lambs and pregnant or lactating does/ewes are more susceptible and need special attention.
- *Haemonchus* is by far the most important cause of anemia in sheep and goats; however, there are other causes of anemia that could cause confusion. Some examples are:
 - Hookworms (very uncommon in US)
 - Live fluke (most likely only a problem in the Gulf Coast and Northwestern States)
 - External parasites
 - Blood parasites (very uncommon in US)
 - Bacterial and viral infections
 - Nutritional deficiencies
- On the other hand, certain conditions can make the eyes' membranes appear redder than expected and thus mask the presence of anemia. Some examples are:
 - Hot and/or dusty conditions that may irritate the eyes
 - Driving animals a long distance with no rest period afterwards

- Fever
- Infectious eye diseases
- Diseases associated with blood circulatory failure
- With FAMACHA[®], animals are allowed to become anemic prior to being treated. Therefore, ***it is critical to use an effective dewormer.*** Drug resistance to all available dewormers is becoming quite common. **Therefore, testing to determine which drugs are effective against the worms on your farm should be done before applying the FAMACHA[®] system.**
 - Available tests for resistant include the fecal egg count reduction tests which are performed on the farm by your veterinarian, and the DrenchRite[®] test, which is performed in a laboratory from a fresh fecal sample this is mailed to the lab.
- Protect the card from light when not in use and replace the card after 1 year of use.

PRACTICAL USE OF THE FAMACHA[®] SYSTEM

- Beginning in the spring or several weeks prior to lambing/kidding, examination of the herd or flock using FAMACHA[®] should be made every 2-3 weeks by properly trained persons.
- During the high worm transmission periods (warm weather), it may be necessary to monitor the flock/herd more often, even on a weekly basis.
- The FAMACHA[®] guide should always be used on inspections. Do not rely on memory from previous examinations.
- Treatment can be safely withheld from adult animals until they score as 4 or 5 if animals are in good body condition and good overall general health, are examined frequently (e.g., every 2-3 weeks) and good husbandry is used to identify animals in need of treatment (e.g., unthrifty, lagging behind, bottle jaw) between FAMACHA[®] examinations. It is advisable to treat animals scored as 3s if any of these conditions are not met.
- Lambs and kids have comparatively small blood volumes and can progress rapidly from moderate to severe anemia. Ewes and does have decreased immunity to worms starting approximately 2 weeks before lambing/kidding and extending through the lactation period (called periparturient period). These animals should always be treated if scored as 3s.
- If 5-10% or more of the herd or flock is found to be anemic (scores of 4 and 5) at any examination, it may be advisable to dose all animals scored in 3-5 and change pastures if available. Such periods of significant *Haemonchus* challenge appear to be heralded by a rapid downward trend in the number of 1s and a reciprocal increase in the number of 2s and 3s. Consult your veterinarian if in doubt.
- The essential decision to be taken at each examination is which animals are to be treated and which are not. Assignment to precise categories is less important. It is better to err on the side of treatment when you are unsure.
- All animals treated with dewormer should be marked or identified in some permanent way (ear tags, ear marks, notches, cable ties, etc.) unless individual animal records are kept. It is recommended that animals that are permanently marked should also be given a temporary mark so that the same animals is not unfairly marked permanently at the next examination.
- Animals needing two doses more than the herd or flock's average should be considered for culling, while those needing three or more extra doses should be culled.
- The proportions of the flock in each category (from 1-5) can easily be recorded by counting off each animal in the FAMACHA[®] block histogram (anemia score sheet) provided.

- If the herd or flock is very large, a random sample of 50 animals can be checked. If the combined percentage of categories 1 and 2 exceeds 80% (preferably 90%) and there are no category 4 and 5 sheep in the sample, it is unlikely that there is danger in not checking the whole flock. However, if any sheep are scored as 4, 5 or the 3 categories exceeds 10-20%, it would be safer to examine all animals.
- Examine especially those animals that lag behind the herd or flock whether or not it's time for a scheduled FAMACHA® examination. These late-comers may be suffering from the effects of anemia.
- Always check animals for "bottle jaw" (presence of soft swelling under the jaw). All animals with bottle jaw, whether they appear anemic or not, must be treated.

FOR INQUIRIES AND ORDER INFORMATION PLEASE CONTACT:

US AND CANADA ONLY

Dr. Adriano Vatta
Louisiana State University
famachasystem@gmail.com
School of Veterinary Medicine
FAMACHA@lsu.edu
(225) 578-9710

INTERNATIONAL

Prof. Gareth Bath
gareth.bath@up.ac.za

COPYRIGHT©

The entire concept, illustrations, and text of this system is subject to copyright rules and no part may be altered or copied in any way without the permission of the copyright holders, the Livestock Health and Production Group of South Africa Veterinary Associates.